# Computational Topology: Isotopic Convergence to a Stick Knot

J. Li<sup>a</sup>, T. J. Peters<sup>1a</sup>, K. E. Jordan<sup>b</sup>, P. Zaffetti<sup>c</sup>

<sup>a</sup>University of Connecticut, Storrs, CT 06269, U.S.A. <sup>b</sup>IBM T.J. Watson Research Center, Cambridge, MA, U.S.A. <sup>c</sup>iDevices Inc., Avon, CT 06001, U.S.A.

# Abstract

Computational topology is a vibrant contemporary subfield and this article integrates knot theory and mathematical visualization. Previous work on computer graphics developed a sequence of smooth knots that were shown to converge point wise to a piecewise linear (PL) approximant. This is extended to isotopic convergence, with that discovery aided by computational experiments. Sufficient conditions to attain isotopic equivalence can be determined *a priori*. These sufficient conditions need not be tight bounds, providing opportunities for further optimizations. The results presented will facilitate further computational experiments on the theory of PL knots (also known as stick knots), where this theory is less mature than for smooth knots.

Key words: Knot, isotopy, computer animation, molecular simulation

## 1. Introduction & Related Work

For a positive integer n, a Bézier curve Piegl and Tiller (1997) of degree n is defined as  $\mathbf{B}(t)$ , with control points  $P_i \in \mathbb{R}^3$  by

$$\mathbf{B}(t) = \sum_{i=0}^{n} \binom{n}{i} t^{i} (1-t)^{n-i} P_{i}, \quad t \in [0,1].$$

The curve  $\mathcal{P}$  formed by PL interpolation on the ordered set of points  $\{P_0, P_1, \ldots, P_n\}$  is called the *control polygon*. This  $\mathcal{P}$  is a PL approximation of **B**.

The curves considered here will be closed by understanding that  $P_0 = P_n$ . Furthermore, it is assumed that both the Bézier curves and their control polygons are simple. The focus here is on the isotopic equivalence between a knotted Bézier curve and its PL approximation. These knotted PL approximations are also known as 'stick knots' Adams et al. (1997).

<sup>&</sup>lt;sup>1</sup>This author (and the 1st author) was partially supported by NSF grant CNS 0923158 and by IBM Faculty Award 2013 - 6327140. All statements here are the responsibility of the author, not of the National Science Foundation nor of IBM.

A contemporary treatment of knots and molecules Wertheim and Millett (2006) provided motivation for much of this work. In particular, remarks on the "... humbling ..." status in theoretical understanding of stick knots versus smooth knots stimulated the consideration, here, of Bézier curves that are isotopically equivalent to stick knots.

The term 'molecular movies' was introduced McGill (27 June 2008) to include the visualization of molecular simulations. There is a specific cautionary example Li et al. (2015) about introducing topological artifacts into a molecular movie. The relevant knots were  $4_1$ and the unknot, represented in sufficiently simple format that isotopic equivalences were easily determined by standard methods Livingston (1993). Knot visualization software Marsh and Peters (2008) was used to develop that illustrative example.

Those application specific considerations Li et al. (2015) led to the generalizations presented here. The theorem presented here generalizes that example to sufficient conditions for convergence between Bézier curves and their isotopic PL approximations. An *a priori* bound is given on the number of iterations needed to obtain an isotopic approximation.

The preservation of topological characteristics in computational applications is of contemporary interest Amenta et al. (2003); Andersson et al. (1995, 1998); Chazal and Cohen-Steiner (2005); Chazal et al. (2011); Denne and Sullivan (2008); Etiene et al. (2012); Jordan et al. (2014b, 2008); Kirby and Silva (2008); Li and Peters (2013); Maekawa et al. (1998). The isotopy theorems on knots of finite total curvature are foundational to the work presented here. Sufficient conditions for a homeomorphism between a Bézier curve and its control polygon have been studied Neagu et al. (2000), while topological differences have also been shown Bisceglio et al. (2011); Li et al. (2012); Piegl and Tiller (1997). Sufficient conditions were given to insure that perturbations of the control points maintain isotopic equivalence of the perturbed splines Andersson et al. (2000). There is an example of a PL structure that becomes self-intersecting while the associated Bézier curve remains simple Cassidy et al. (2014). Recent perspectives on computational topology have appeared Carlsson (2009); Edelsbrunner and Harer (2010); Zomorodian (2005).

# 2. Inserting Midpoints as Control Points

The fundamental approximation technique introduced here is dual to many others. The more typical focus is to produce a sequence of PL curves that approximates a given smooth curve. Indeed, these authors have previously published such results Jordan et al. (2014a). The duality here is to create a sequence of smooth knots that converge to a PL knot, achieving isotopic equivalence within the sequence. The use of smoothness here is understood to be  $C^{\infty}$ , with a possible exception at the point  $\mathbf{B}(0) = \mathbf{B}(1)$ . This technique has been called *collinear insertion* Li et al. (2015), where an example was presented with 8 initial control points (inclusive of equality of the initial and final control points). That control polygon was the knot  $4_1$  and its associated Bézier curve was the unknot. With 4 iterations of collinear insertion, the stick and smooth knots were isotopic.

**Definition 2.1.** Consider the control polygon  $\mathcal{P}$ . To avoid trivial cases, it is assumed that, for i > 0,

- $P_i \neq P_i *$  for any  $i \neq i *$ ,
- only two such  $P_i \neq P_i *$  can be collinear and
- $n \ge 4$ .

Sequences of control polygons and Bézier curves will be generated by letting  $\mathcal{P}^{(0)} = \mathcal{P}$  and generating  $\mathcal{P}^{(1)}$  from  $\mathcal{P}^{(0)}$  by adding the midpoint of each edge of  $\mathcal{P}^{(0)}$ . For  $j \geq 0$ , similarly generate  $\mathcal{P}^{(j+1)}$  by the insertion of midpoints between all of the control points of  $\mathcal{P}^{(j)}$ . For each j, the corresponding Bézier curve will be denoted as  $\mathbf{B}^{(j)}$ . Note that all  $\mathcal{P}^{(j)}$ 's are isotopic under the trivial identification map.

# 3. Convergence Theorem

The primary convergence result relies upon a previously published theorem (Denne and Sullivan, 2008, Theorem 4.2) on rectifiable Morgan (2008) graphs of finite total curvature Milnor (1950). This central theorem is quoted, below, after definition of a key notion of closeness.

**Definition 3.1.** Given two rectifiable embeddings  $\Gamma_1$  and  $\Gamma_2$  of the same combinatorial graph, we say they are  $(\delta, \theta)$  – close if there exists a homeomorphism between them such that corresponding points are within distance  $\delta$  of each other, and corresponding tangent vectors are within angle  $\theta$  of each other almost everywhere.

**Theorem 3.1.** (Denne and Sullivan, 2008, Theorem 4.2) Suppose  $\Gamma_1$  is a knotted graph of finite total curvature and  $\epsilon > 0$  is given. Then there exists  $\delta > 0$  such that any (rectifiable) graph  $\Gamma_2$  which is  $(\delta, \pi/8)$  – close to  $\Gamma_1$  is ambient isotopic to  $\Gamma_1$ , via an isotopy which moves no point by more than  $\epsilon$ .

The tangent vectors of the Bézier curves and their control polygons are well defined except at possibly finitely many points. The homeomorphism between the knotted control polygon and knotted Bézier curve will be the natural one matching points defined at the same parametric values over  $[0, 2\pi]$ . The PL knots considered will have finite total curvature. The result that every compact connected set which has finite length is rectifiable David and Semmes (1993) is directly applicable to the curves considered.

There is pervasive interest in computer graphics in convergence properties between Bézier curves and their PL approximations Piegl and Tiller (1997), motivating the general convergence considered here for collinear insertion. There are supportive techniques already available for Bézier curves, as summarized briefly here.

The derivative of **B** is a Bézier curve Farin (1990), with control points  $n * \Delta P_i$ , in terms of the forward difference operator, denoted by

$$\Delta P_i = P_{i+1} - P_i, \ i = 0, \dots, n - 1.$$

The second iterated forward difference operator is defined similarly as

$$\Delta_2 P_i = P_{i+2} - 2P_{i+1} + P_i.$$

Point wise convergence in distance under j collinear insertions has been shown Li et al. (2015). Bounds will be shown on the rate of convergence for collinear insertion by adapting a previously published inequality Nairn et al. (1999). Each control point of **B** is an ordered triple  $(x, y, z) \in \mathbf{R}^3$ . Let  $\Delta_2 P^x$  denote the second iterated forward difference taken over the x coordinate (with similarly definitions regarding the y and z coordinates). Then,  $\Delta_2 P^x$  is a vector having n-2 entries and the usual 1-norm is indicated by  $\|\Delta_2 P^x\|_1$ . Let

$$\|\Delta_2 P\|_{1,M} = \max_{w \in \{x,y,z\}} \|\Delta_2 P^w\|_1.$$

The notation Nairn et al. (1999) of  $N_1(n)$  and a related combinatorial upper bound appears in the appendix of Section 6. The indicated adaptation was previously implicit Li et al. (2015)but is explicated as

$$\|\mathbf{B}(t) - \mathcal{P}(t)\|_{\infty,[0,1]} \le N_1(n) * \|\Delta_2 P\|_{1,M},$$
(3.1)

leading directly to

$$\|\mathbf{B}^{(j)}(t) - \mathcal{P}^{(j)}(t)\|_{\infty,[0,1]} \le N_1(2^j n) * \|\Delta_2 P^{(j)}\|_{1,M} \le (n/(4\sqrt{n * 2^j + 1}))\|\Delta_2 P\|_{1,M}.$$
 (3.2)

#### 3.1. Distance Bounds: Curve and Hodograph

An upper bound on the distance between a hodograph and its control polygon is also determined directly from Inequality 3.1. Let  $\frac{d}{dt}B$  denote the derivative of **B** and  $\frac{d}{dt}B$  is a Bézier curve with control points of the form

$$n * (P_i - P_{i-1}), \quad i = 1, \dots, n-1.$$
 (3.3)

Let  $H_1, H_2, \ldots, H_{n-1}$  be the control points of  $\frac{d}{dt}\mathbf{B}$  and let  $\mathcal{H}$  be the associated control polygon, leading to the inequality

$$\|\frac{\mathbf{d}}{\mathbf{dt}}\mathbf{B}(t) - \mathcal{H}(t)\|_{\infty,[0,1]} \le N_1(n) * \|\Delta_2 H\|_{1,M}.$$
(3.4)

A slight refinement has been made on the right hand side of Inequality 3.4 in the use of  $N_1(n)$  in place of  $N_1(n-1)$ , as  $N_1$  is a " ... slowly growing function ..." Nairn et al. (1999). The advantage of this minor replacement is to continue to evaluate  $N_1$  on n as an even integer, where that convenience is justified in Section 6. While this increases the upper bound, the increase is not substantive for theoretical convergence questions, though a tighter upper bound may be valuable to consider on specific data.

For each  $j \in \mathbf{N} \cup \{0\}$ , let  $\mathcal{H}^{(j)}$  denote the control polygon for  $\frac{\mathbf{d}}{\mathbf{dt}} \mathbf{B}^{(j)}(t)$ . By definition,

$$\Delta_2 H_{i+2} = n * ((P_{i+2} - P_{i+1}) - 2(P_{i+1} - P_i) + (P_{i+2} - P_{i-1})),$$

which simplifies to

$$\Delta_2 H_{i+2} = n * (P_{i+2} - 3P_{i+1} + 3P_i - P_{i-1}).$$
(3.5)

Lemma 3.1. For each  $j \in \mathbf{N} \cup \{0\}, \|\Delta_2 H^{(j)}\|_{1,M} \le 2^{j-1} \|\Delta_2 H\|_{1,M}.$ 

**Proof:** For each j, there are two important cases to consider for  $\Delta_2 H^{(j)}$ . Recall that the distance between successive control points of  $P^{(j)}$ , is equal  $2^{-j} ||P_{\ell} - P_{\ell-1}||$ , for  $\ell = 1, \ldots n$ .

Case 1: The four points used in Equation 3.5 are collinear and  $\Delta_2 H_i^{(j)}$  is zero over all coordinates x, y, z.

Case 2: Three of the four points in Equation 3.5 are collinear. If the first three points are collinear, then  $\|\Delta_2 H_i^{(j)}\| = 2^{-j} * n \|P_{\ell} - P_{\ell-1}\|$  for some  $\ell = 1, \ldots n$ . If the last three points are collinear, then the same value is obtained. Hence, in evaluation of the 1-norm for the vector  $\Delta_2 \mathcal{H}^{(j)}$ , this term occurs twice for each  $\ell$ .

The conclusion from these two cases is that

$$\|\Delta_2 \mathcal{H}^{(j)}\| = 2^{-(j-1)} \|\Delta_2 \mathcal{H}\|. \qquad \Box$$

Applying Lemma 3.1 and Inequality 3.2 to the hodograph yields

$$\|\frac{\mathbf{d}}{\mathbf{dt}}\mathbf{B}^{(j)}(t) - \mathcal{H}^{(j)}(t)\|_{\infty,[0,1]} \le \left(n/(2\sqrt{n*2^{j}+1})\right)*2^{-(j-1)}*\|\Delta_{2}H\|_{1,M}.$$
 (3.6)

#### 3.2. Bounding Angles Between Tangent Vectors

Inequality 3.6 provides a bound on distance between each point on the hodograph,  $\frac{d}{dt}\mathbf{B}^{(j)}(t)$  and the corresponding point  $\mathcal{H}^{(j)}(t)$  on the control polygon, which suffices as a bound between corresponding tangent vectors.

For each non-negative integer j, there are  $2^{j}n + 1$  control points of  $\mathcal{P}^{(j)}$ . Let  $k = 0, \ldots, 2^{j}n$ and define a uniform parametrization of [0,1] with  $\mathcal{P}^{(j)}(k/(2^{j}n))$  being the k-th control point of  $\mathcal{P}^{(j)}$  and other points on the control polygon determined by linear interpolation. Closed parametric subintervals are of the form  $[(k/(2^{j}n), ((k+1)/(2^{j}n)]]$ . The presence of consecutive collinear control points has implications for the discrete derivative.

In Figure 1, the triangle depicts the subtraction of any two vectors, where upper case letters denote angles, while lower case letters denote edges. Consider **A** relative to an upper bound on  $\|\mathbf{a}\|$ , where the direction of the vector corresponding to edge  $\|\mathbf{a}\|$  is unknown. The dotted circle indicates the positions that this edge could assume. By symmetry, only the semicircle above the horizontal axis needs to be considered. When  $\|\mathbf{a}\| < \|\mathbf{b}\|$ , angle **A** attains its maximum when angle **C** is  $\pi/2$ , by elementary trigonometry.

**Lemma 3.2.** The maximum angle between tangent vectors of  $\mathbf{B}^{(j)}$  and  $\mathcal{P}$  can be determined over the shortest edge of  $\mathcal{P}$ .

**Proof:** For each  $t \in [0, 1]$ , consider  $\frac{\mathbf{d}}{\mathbf{dt}} \mathbf{B}^{(j)}(t) - \mathcal{H}^{(j)}(t)$ , with edge **b** in Figure 1 depicting the tangent of  $\mathcal{P}$  that corresponds to  $\frac{\mathbf{d}}{\mathbf{dt}} \mathbf{B}^{(j)}(t)$ .

If t is in an interval  $[(k/(2^{j}n), ((k+1)/(2^{j}n)]]$  which does not contain some i/n for any i = 0, ..., n, then  $\mathbf{b} = n ||P_{i+1} - P_i||$ , for some i = 0, ..., n - 1. In the notation of Figure 1, the maximum value for angle  $\mathbf{A}$  will be given by  $\arcsin(\mathbf{a}/\mathbf{b})$ .



Figure 1: Upper Bound on Angle

Denote by  $\lambda$  the minimum length over all edges from  $\mathcal{P}$  and let **b** be a tangent vector of the form  $n * (P_i - P_{i-1})$ , for some  $i = 1, \ldots, n$ . The condition on the length of **a** is met by choosing j such that the right hand side of Inequality 3.6 is less than  $n\lambda$ .  $\Box$ 

#### 4. An Example

Following from Lemma 3.2 and Inequality 3.6 the values to satisfy Theorem 3.1 can be determined with an integer  $m_1$  sufficient to achieve the distance bound of  $\delta$  and with an integer  $m_2$  sufficient to achieve the angular bound of  $\pi/8$ . Let  $M = max\{m_1, m_2\}$ , so that for all j > M,  $\mathbf{B}^{(j)}$  is isotopic to  $\mathcal{P}$ .

A detailed example follows on the control polygon Li et al. (2015)

$$(1.3076, -3.3320, -2.5072), (-1.3841, 4.6826, 0.9135), (-3.2983, -4.0567, 2.6862), (-0.1233, 2.7683, -2.4636), (3.9080, -4.5334, 1.2264), (-3.9360, -0.4383, -0.9834), (3.2182, 4.2961, 2.1125).$$

#### 4.1. Empirically Determined Bounds

Figure 2 depicts the stick knot  $4_1$ , successively followed by the Bézier curves of the unknot from the 0th collinear insertion and of  $4_1$  from the 4th collinear insertion, as previously proven Li et al. (2015). Figure 3 shows Iterations 1 - 3 of collinear insertion under projections that provide visual verification that each of the corresponding Bézier curves is the unknot.

#### 4.2. Theoretically Determined Bounds

To apply Theorem 3.1 note that  $\mathcal{P}$  has finite total curvature.

The following narrative to compute  $\delta$  is adapted from the source Denne and Sullivan (2008) for the convenience of the reader. Select a finite number of points  $p_j$  from  $\mathcal{P}$  (including all its control points) such that these points divide  $\mathcal{P}$  into PL arcs  $\alpha_k$ , each of total curvature less than  $\pi/8$ . Let  $\epsilon = 1.0$ 



Figure 3: Projections of Three Iterations of Collinear Insertion.

Let  $r_1$  be the minimum distance between any two arcs  $a_k$  which are not incident to a common  $p_j$  (or the minimum distance between points  $p_j$ , if this is smaller). Let

$$r_2 = \min(r_1/2, \epsilon/2).$$

Consider disjoint open balls  $B_j$  of radius  $r_2$  centered at the  $p_j$ . Note that  $\mathcal{P} \setminus \bigcup B_j$  is a compact union of disjoint arcs  $\beta_k \subset \alpha_k$ . Let  $r_3$  be the minimum distance between any two of these arcs  $\beta_k$ 

Let  $r_4 = r_3/6$  and let  $\delta = r_4/3$ .

The values for  $\mathcal{P}$  are summarized in Table 1.

Variable	Value
$\epsilon$	1.0000
$r_1$	0.2576
$r_2$	0.1288
$r_3$	0.0576
$r_4$	0.0096
δ	0.0032

Table 1: Computing  $\delta$ .

The values for  $m_1$  and  $m_2$  discussed at the beginning of this section can now be calculated, using  $n = 7, \lambda = 9$ .

To obtain  $m_1$ , use Inequality 3.2 and let  $\delta = (n/(4\sqrt{n*2^j+1}))* \|\Delta_2 P\|_{1,M}$ . Let  $\Omega_1 = \|\Delta_2 P\|_{1,M} = 70.8$ , so that

$$m_1 = \lceil \log_2((7\Omega_1^2)/(16\delta^2) - (1/7)) \rceil = 28.$$

There are two constraints on  $m_2$  in Lemma 3.2, with the first being relative to the shortest tangent vector and the second in terms of the angular bound of  $\pi/8$ . The corresponding number of iterations to meet those constraints will be denoted by  $m_{2T}$  and  $m_{2A}$ , respectively, with  $m_2 = \max\{m_{2T}, m_{2A}\}$ .

First,  $m_{2T}$  is computed following Lemma 3.2 to have  $||a|| < n\lambda$ , with ||a|| set equal to the right hand side (RHS) of Inequality3.6 and  $\Omega_2 = ||\Delta_2 H||_{1,M} = 779.0$  to get

$$(n/(2*\sqrt{n*2^{j}})*2^{-(j-1)}*\Omega_{2} < n\lambda,$$

and

$$(\Omega_2/\lambda)^2 < n2^{3j} + 2^{2j},$$

to yield  $m_{2T} = 4$ .

For  $m_{2A}$ , again, set ||a|| equal to the RHS of Inequality3.6 and use  $\arcsin(||a||/(n\lambda)) = \pi/8$  to get  $m_{2A} = 5$ . It is then clear, that  $m_2 = 5$  and M = 28.

The difference between the theoretically determined sufficient condition of 28 iterations to achieve isotopic equivalence versus the visual inspection that 4 iterations suffice present opportunities to explore tightness criteria. These future investigations will be enabled by previously developed knot visualization software Marsh and Peters (2008). It is of interest to note that the empirically determined optimal value of 4 iterations for the example is exactly met by the tangency condition, with  $m_{2T} = 4$ , and is nearly met by the angular condition, with  $m_{2A} = 5$ . This directs initial attention to considering why the distance criteria yield such a larger number of 28, while also appreciating that the tangency and angular conditions for this example might just be particularly well-behaved.

#### 5. Conclusion and Future Work

For synchronous visualizations of writhing molecules, a cautionary example was previously presented of different knot types between a polynomial curve and its rendering. Further study of the isotopy constraints on that example prompted these generalizations for a sequence Bézier curves converging to a given stick knot. This broader theory will further inform the design of visualization software for molecular simulations. Beyond that application, this work lays the foundation for mathematical visualization software for the experimental investigation of important theoretical relationships between stick and smooth knots, where much remains to be discovered. A particular future emphasis will be upon whether isotopic approximations can occur earlier in the converging sequence than given by the sufficient *a prior* bounds presented here. Any discovery of more aggressive bounds will likely rely upon computational experiments to discover criteria for tight bounds.

### 6. Appendix: Supportive Combinatorial Relations

Let N denote the natural numbers  $\{1, 2, 3, ...\}$ . For  $k \in \mathbb{N}$ , consider the central binomial coefficient Chowdhury et al. (2014),

$$\left(\begin{array}{c}2k\\k\end{array}\right).$$

Lemma 6.1. For  $k \in \mathbf{N}$ ,

$$\left(\begin{array}{c}2k\\k\end{array}\right) \le \frac{4^k}{\sqrt{2k+1}}.$$

**Proof:** The key to the proof has informally been attributed to P. Erdos Chowdhury et al. (2014).

The following notation has previously appeared Nairn et al. (1999) and is central to the convergence results. Note that presenting the definition purely for even arguments is sufficient here, since the colliear insertion process always doubles the degree of the previous Bézier curve. For  $k \in N$ , let

$$N_1(2k) = \begin{pmatrix} 2k \\ k \end{pmatrix} \frac{2k}{2^{2k+2}}.$$

Lemma 6.2. For  $k \in N$ ,

$$N_1(2k) < \frac{k}{2\sqrt{2k+1}}.$$

**Proof:** The short proof Li et al. (2015) is repeated here for the convenience of the reader. Invoke Lemma 6.1 on

$$N_1(2k) = \binom{2k}{k} \frac{2k}{2^{2k+2}} < \frac{4^k}{\sqrt{2k+1}} \frac{2k}{4^{k+1}} = \frac{k}{2\sqrt{2k+1}}. \qquad \Box$$

# References

- Adams, C. C., Brennan, B. M., Greilsheimer, D. L., Woo, A. K., 1997. Stick numbers and composition of knots and links. Journal of Knot Theory and its Ramifications 6 (02), 149–161.
- Amenta, N., Peters, T. J., Russell, A. C., 2003. Computational topology: Ambient isotopic approximation of 2-manifolds. Theoretical Computer Science 305, 3–15.
- Andersson, L. E., Dorney, S. M., Peters, T. J., Stewart, N. F., 1995. Polyhedral perturbations that preserve topological form. CAGD 12(8), 785–799.
- Andersson, L. E., Peters, T. J., Stewart, N. F., 1998. Selfintersection of composite curves and surfaces. CAGD 15, 507–527.
- Andersson, L.-E., Peters, T. J., Stewart, N. F., 2000. Equivalence of topological form for curvilinear geometric objects. International Journal of Computational Geometry & Applications 10 (06), 609–622.
- Bisceglio, J., Peters, T. J., Roulier, J. A., Sequin, C. H., 2011. Unknots with highly knotted control polygons. CAGD 28(3), 212–214.
- Carlsson, G., 2009. Topology and data. Bulletin of the American Mathematical Society 46 (2), 255–308.
- Cassidy, H. P., Peters, T. J., Ilies, H., Jordan, K. E., 2014. Topological integrity for dynamic spline models during visualization of big data. Topological Methods in Data Analysis and Visualization III, 110–121.
- Chazal, F., Cohen-Steiner, D., 2005. A condition for isotopic approximation. Graphical Models 67 (5), 390–404.
- Chazal, F., Cohen-Steiner, D., Mérigot, Q., 2011. Geometric inference for probability measures. Foundations of Computational Mathematics 11, 733–751.
- Chowdhury, M. B. R., Komjth, P., Speyer, D., 2014. Upper limit on the central binomial coefficient. http://mathoverflow.net/questions/133732/upper-limit-on-the-central-binomial-coefficient.
- David, G., Semmes, S., 1993. Analysis of and on uniformly rectifiable sets. Vol. 38. American Mathematical Soc.
- Denne, E., Sullivan, J. M., 2008. Convergence and isotopy type for graphs of finite total curvature. In: Bobenko, A. I., Sullivan, J. M., Schröder, P., Ziegler, G. M. (Eds.), Discrete Differential Geometry. Birkhäuser Basel, pp. 163–174.
- Edelsbrunner, H., Harer, J., 2010. Computational topology: an introduction. American Mathematical Soc.

- Etiene, T., Nonato, L. G., Scheidegger, C. E., Tierny, J., Peters, T. J., Pascucci, V., Kirby, R. M., Silva, C. T., 2012. Topology verification for isosurface extraction. IEEE Trans. Vis. Comput. Graph. 18 (6), 952–965.
- Farin, G. E., 1990. Curves and Surfaces for Computer-Aided Geometric Design: A Practical Guide. Academic Press, Inc.
- Jordan, K., Li, J., Peters, T., Roulier, J., 2014a. Isotopic equivalence by bézier curve subdivision for application to high performance computing. Computer Aided Geometric Design 31 (9), 642–655.
- Jordan, K. E., Li, J., Peters, T. J., Roulier, J. A., 2014b. Isotopic equivalence from Bézier curve subdivision for application to high performance computing. CAGD 31, 642 655.
- Jordan, K. E., Miller, L. E., Moore, E. L. F., Peters, T. J., Russell, A., 2008. Modeling time and topology for animation and visualization with examples on parametric geometry. Theoretical Computer Science 405, 41–49.
- Kirby, R. M., Silva, C. T., 2008. The need for verifiable visualization. IEEE Computer Graphics and Applications September/October, 1–9.
- Li, J., Peters, T., Marinelli, K., Kovalev, E., Jordan, K., 2015. Topological subtleties for molecular movies. Topology and its Applications 188, 91–96.
- Li, J., Peters, T. J., 2013. Isotopic convergence theorem. Journal of Knot Theory and Its Ramifications 22 (3).
- Li, J., Peters, T. J., Marsh, D., Jordan, K. E., 2012. Computational topology counterexamples with 3D visualization of Bézier curves. Applied General Topology 13 (2), 115–134.
- Livingston, C., 1993. Knot Theory. Vol. 24 of Carus Mathematical Monographs. Mathematical Association of America, Washington, DC.
- Maekawa, T., Patrikalakis, N. M., Sakkalis, T., Yu, G., 1998. Analysis and applications of pipe surfaces. CAGD 15(5), 437–458.
- Marsh, D. D., Peters, T. J., 2008. Knot and bezier curve visualizing tool. http://www.cse.uconn.edu/~tpeters/top-viz.html.
- McGill, G., 27 June 2008. Molecular movies coming to a lecture near you. Cell 133 (7), 1127–1132.
- Milnor, J. W., 1950. On the total curvature of knots. Annals of Mathematics 52, 248–257.
- Morgan, F., 2008. Geometric measure theory: a beginner's guide. Academic press.
- Nairn, D., Peters, J., Lutterkort, D., 1999. Sharp, quantitative bounds on the distance between a polynomial piece and its Bézier control polygon. CAGD 16, 613–631.

Neagu, M., Calcoen, E., Lacolle, B., 2000. Bézier curves: topological convergence of the control polygon. 6th Int. Conf. on Mathematical Methods for Curves and Surfaces, Vanderbilt, 347–354.

Piegl, L., Tiller, W., 1997. The NURBS Book. Springer, New York.

Wertheim, M., Millett, K., 2006. Where the wild things are: An interview with Ken Millett. Cabinet 20.

Zomorodian, A. J., 2005. Topology for computing. Vol. 16. Cambridge University Press.